

# STAT 315: Binomial Random Variables

Luc Rey-Bellet

University of Massachusetts Amherst

*luc@math.umass.edu*

September 29, 2025

# Binomial trials

## Binomial experiments

- The experiment consists of  $n$  identical trials.
- Each trial results in one of two outcomes: success  $S$  or failure  $F$ .
- Each trial has probability of success  $p$  and probability of failure  $q = (1 - p)$ .
- The trials are independent.
- The binomial random variable  $Y$  is

$Y$  = number of successes observed during the  $n$  trials

The random variable  $Y$  has two parameters

$n$  = numbers of trials

$p$  = probability of success

**Notation:** We write  $Y \sim B_{n,p}$

# Math reminder: The binomial theorem

## Theorem

*The binomial theorem*

$$(x + y)^n = \sum_{k=0}^n \binom{n}{k} x^k y^{n-k}$$

**Proof:**

$$(x + y)^n = \underbrace{(x + y)(x + y) \cdots (x + y)}_{n \text{ times}}$$

Expanding the product gives a sum of  $2^n$  terms. Each term has the form  $x^k y^{n-k}$  with  $k = 0, 1, \dots, n$ .

The number of terms of the form  $x^k y^{n-k}$  is equal to  $\binom{n}{k}$  since it is the number of ways to choose the  $k$  locations for the  $x$ .

Adding all the terms up give the theorem.

# The PDF of a binomial random variable

## Probability distribution function

- $Y$  takes values  $0, 1, 2, \dots, n$  (the number of successes).
- The pdf is

$$p(k) = P(Y = k) = \binom{n}{k} p^k (1 - p)^{n-k}$$


For example with  $n = 5$  a sequence of 2 failures and three successes e.g. *FSSFS* has probability  $p^3(1 - p)^2$  and there are  $\binom{5}{3}$  such sequences (=number of ways to choose which ones of the five trials is a success).

Note that by the binomial theorem

$$1 = 1^n = (p + (1 - p))^n = \sum_{k=0}^n \binom{n}{k} p^k (1 - p)^{n-k} = \sum_{k=0}^n p(k)$$

and so  $p(k)$  is a pdf!

# Shape of the binomial distribution



**Figure:** The binomial pdf for  $B_{20,.04}$ ,  $B_{20,.16}$ , and  $B_{20,.53}$  from left to right.

$$\frac{p(k)}{p(k-1)} = \frac{\frac{n!}{(k)!(n-k)!} p^k (1-p)^{n-k}}{\frac{n!}{(k-1)!(n-k+1)!} p^{k-1} (1-p)^{n-k+1}} = \frac{n-k+1}{k} \frac{p}{1-p}$$

and thus  $\frac{p(k)}{p(k-1)} \geq 1$  iff  $(n-k+1)p > k(1-p)$  iff  $(n+1)p \geq k$

The maximum is around  $k_{max} \approx (n+1)p$ .

# Applications of the Binomial Distribution

- **Quality Control & Reliability:** number of defective items in a batch.
- **Clinical Trials:** number of patients cured with a given probability.
- **Elections & Polling:** number of voters supporting a candidate in a sample.
- **Insurance & Risk:** number of claims or accidents among insureds.
- **Genetics:** inheritance of a trait across offspring.
- **Communication Systems:** number of corrupted bits in a data packet.
- **Gambling & Games:** number of heads in coin flips or wins in repeated trials.

## Example


- A multiple-choice examination has 10 questions, each with four possible answers, only one of which is correct. Having not studied a student answers each of the questions with an independent random guess.
  - ▶ What is the probability he get 2 or 3 questions correct?
  - ▶ What is the probability the students gets a passing score (at least 6 section correct)?
- Suppose each child of two carriers of a recessive gene has a 25% chance of being affected. If a couple has  $n = 4$  children, let  $X =$  number of affected children.
  - ▶ What is the probability that exactly 2 children are affected?
  - ▶ What is the probability that at least one child is affected?

# Binomial Distribution: $n = 10, p = 1/4$

$$X \sim \text{Binomial}(10, 1/4)$$

## Probabilities:

- $P(X = 0) \approx 0.0563$
- $P(X = 1) \approx 0.1877$
- $P(X = 2) \approx 0.2817$
- $P(X = 3) \approx 0.2500$
- $P(X = 4) \approx 0.1406$
- $P(X = 5) \approx 0.0540$
- $P(X = 6) \approx 0.0162$
- $P(X = 7) \approx 0.0031$
- $P(X = 8) \approx 0.00039$
- $P(X = 9) \approx 0.000029$
- $P(X = 10) \approx 0.000001$





# Chuck-a-luck (USA) or Bầu cua tôm cá (gourd-crab-shrimp-fish, Vietnam) or Hoo Hey How (Fish-Prawn-Crab, China)



The game is played with 3 six-faced dice a board with the pictures (or numbers) on the six faces.

If you bet \$1 on say "shrimp", you win \$ $k$  if you land  $k$  "shrimps",  $k = 1, 2, 3$  and lose your \$1 otherwise. Find the pdf, expected value and variance of the gain.

# Mean and Variance of the binomial RV

## Mean and Variance of a binomial RV $Y \sim B_{n,p}$

If  $n$  is the number of trials and  $p$  the probability of success

$$E[Y] = \mu = np$$

$$V[Y] = \sigma^2 = np(1 - p)$$

Note that  $E[Y]$  is order  $n$  and so maybe one would guess that  $V[Y] = E[Y^2] - E[Y]^2$  should be of order  $n^2$ . But this is not so (some cancellation occurs..)

Note the following identity

$$k \binom{n}{k} = \frac{kn!}{k!(n-k)!} = \frac{n!}{(k-1)!(n-k)!} = \frac{n(n-1)!}{(k-1)!(n-k)!} = n \binom{n-1}{k-1}$$

Pick a team of  $k$  then a captain versus Pick a captain and then the rest of the team.

$$\begin{aligned} E[Y] &= \sum_{k=0}^n k \binom{n}{k} p^k (1-p)^{n-k} = \sum_{k=1}^n n \binom{n-1}{k-1} p^k (1-p)^{n-k} \\ &= np \sum_{k=1}^n \binom{n-1}{k-1} p^{k-1} (1-p)^{n-k} = np \underbrace{\sum_{j=0}^{n-1} \binom{n-1}{j} p^j (1-p)^{n-1-j}}_{=1} \\ &= np \end{aligned}$$

Similarly

$$k(k-1)\binom{n}{k} = \frac{n!}{(k-2)!(n-k)!} = n(n-1)\binom{n-2}{k-2}$$

and

$$\begin{aligned} E[Y(Y-1)] &= \sum_{k=0}^n k(k-1)\binom{n}{k} p^k (1-p)^{n-k} \\ &= n(n-1)p^2 \sum_{k=2}^n \binom{n-2}{k-2} p^{k-2} (1-p)^{n-k} \\ &= n(n-1)p^2 \underbrace{\sum_{j=0}^{n-2} \binom{n-2}{j} p^j (1-p)^{n-2-j}}_{=1} = n(n-1)p^2 \end{aligned}$$

So

$$\begin{aligned} V(Y) &= E[Y^2] - E[Y]^2 = E[Y(Y-1)] + E[Y] - E[Y]^2 \\ &= n(n-1)p^2 + np - n^2p^2 = np - np^2 = np(1-p) \end{aligned}$$

# Examples

- A multiple-choice examination has 10 questions, each with four possible answers, only one of which is correct. Having not studied a student answers each of the questions with an independent random guess.
  - ▶ What is the mean and variance of the number of correct answers?
- Suppose each child of two carriers of a recessive gene has a 25% chance of being affected. If a couple has  $n = 4$  children, let  $X =$  number of affected children.
  - ▶ What is the mean and variance of the number of affected children.