

STAT 315: Functions of Random Variables I: CDF Method

Luc Rey-Bellet

University of Massachusetts Amherst

luc@math.umass.edu

April 21, 2025

Function of Random Variables

If we have a function of random variables, say

$$Z = g(Y)$$

or

$$Z = g(Y_1, Y_2)$$

or maybe a sample average

$$Z = \frac{1}{n} \sum_{i=1}^n Y_i = \frac{Y_1 + \cdots + Y_n}{n}$$

We know how to compute expectations $E[g(Y)]$ or $E[g(Y_1, Y_2)]$ but often we need more

How do we compute the pdf or cdf of Z ?

The CDF method

The CDF method

For $Z = g(Y_1, Y_2, \dots, Y_n)$ compute the CDF of Z by

- Identify the region $Z = g(Y_1, \dots, Y_n) = z$ in the $y_1, \dots, y_n$ space.
- Identify the region $Z = g(Y_1, \dots, Y_n) \leq z$ in the $y_1, \dots, y_n$ space.
- Compute the integral

$$F(z) = P(Z \leq z) = \int \cdots \int_{g(y_1, \dots, y_n) \leq z} f(y_1, y_2, \dots, y_n) dy_1 dy_2 \cdots dy_n$$

- Compute the pdf of Z by

$$f(z) = F'(z).$$

Example: Linear transformations

Suppose that the random variable X has PDF $f_X(x)$ and CDF $F_X(x)$. Find the density of $Y = aX + b$?

First take $a > 0$:

$$\begin{aligned}\text{CDF of } Y : \quad F_Y(y) &= P(Y \leq y) = P(aX + b \leq y) \\ &= P(aX \leq y - b) \underbrace{=}_{\text{use } a > 0} P\left(X \leq \frac{y - b}{a}\right) \\ &= F_X\left(\frac{y - b}{a}\right).\end{aligned}$$

Differentiating we find

$$f_Y(y) = F'_Y(y) = \frac{d}{dy} F_X\left(\frac{y - b}{a}\right) = F'_X\left(\frac{y - b}{a}\right) \frac{1}{a} = \frac{1}{a} f_X\left(\frac{y - b}{a}\right)$$

Next assume $a < 0$:

$$\begin{aligned}\text{CDF of } Y : \quad F_Y(y) &= P(Y \leq y) = P(aX + b \leq y) \\ &= P(aX \leq y - b) \underbrace{=}_{\text{use } a < 0} P\left(X \geq \frac{y - b}{a}\right) \\ &= 1 - F_X\left(\frac{y - b}{a}\right).\end{aligned}$$

Then

$$f_Y(y) = \frac{d}{dy} \left(1 - F_X\left(\frac{y - b}{a}\right) \right) = -F'_X\left(\frac{y - b}{a}\right) \frac{1}{a} = \frac{1}{-a} f_X\left(\frac{y - b}{a}\right)$$

PDF of $Y = aX + b$

$$X \text{ has PDF } f_X(x) \implies Y = aX + b \text{ has PDF } f_Y(y) = \frac{1}{|a|} f_X\left(\frac{y - b}{a}\right)$$

Example 1: X is a normal RV with mean μ and variance σ^2

$$f_X(x) = \frac{1}{\sigma\sqrt{2\pi}} e^{-\frac{(x-\mu)^2}{2\sigma^2}}$$

Then $Y = aX + b$ has density

$$f_Y(y) = \frac{1}{|a|} f_X\left(\frac{y-b}{a}\right) = \frac{1}{|a|} \frac{1}{\sigma\sqrt{2\pi}} e^{-\frac{\left(\frac{y-b}{a} - \mu\right)^2}{2\sigma^2}} = \frac{1}{|a|\sigma\sqrt{2\pi}} e^{-\frac{(y-(a\mu+b))^2}{2a^2\sigma^2}}$$

so Y is normal with mean $a\mu + b$ and variance $a^2\sigma^2$.

Example 2: Suppose X is an exponential random variable with parameter β . So $f(x) = \frac{1}{\beta} e^{-\frac{x}{\beta}}$

Then $Y = aX$ (with $a > 0$) has density

$$f_Y(y) = \frac{1}{a} f_X(y/a) = \frac{1}{a} \frac{1}{\beta} e^{-\frac{y/a}{\beta}} = \frac{1}{a\beta} e^{-\frac{y}{a\beta}}$$

so Y is exponential with parameter $a\beta$.

The function $Y = X^2$

Given the PDF $f(x)$ of X we want the pdf of $Y = X^2$. We have

$$\{Y \leq y\} \iff \{X^2 \leq y\} \iff \{-\sqrt{y} \leq X \leq \sqrt{y}\}$$

$$\begin{aligned} F_Y(y) &= P(Y \leq y) = P(X^2 \leq y) \\ &= P(-\sqrt{y} \leq X \leq \sqrt{y}) = F_X(\sqrt{y}) - F_X(-\sqrt{y}) \end{aligned}$$

$$\begin{aligned} f_Y(y) &= \frac{d}{dy} F_Y(y) = \frac{d}{dy} (F_X(\sqrt{y}) - F_X(-\sqrt{y})) \\ &= f_X(\sqrt{y}) \frac{1}{2\sqrt{y}} + f_X(-\sqrt{y}) \frac{1}{2\sqrt{y}} \end{aligned}$$

PDF of $Y = X^2$

X has PDF $f_X(x) \Rightarrow Y = X^2$ has PDF $f_Y(y) = \frac{1}{2\sqrt{y}}(f_X(\sqrt{y}) + f_X(-\sqrt{y}))$

The relation between normal RV and χ^2 RV (a.k.a Gamma with $\alpha = \frac{1}{2}$ and $\beta = 2$)

If Z is a standard normal RV, (PDF $f(z) = \frac{1}{\sqrt{2\pi}} e^{-\frac{z^2}{2}}$) then $Y = Z^2$ has pdf

$$\begin{aligned} f_Y(y) &= f_Z(\sqrt{y}) \frac{1}{2\sqrt{y}} + f_Z(-\sqrt{y}) \frac{1}{2\sqrt{y}} \\ &= \frac{1}{\sqrt{2\pi}} e^{-\frac{(\sqrt{y})^2}{2}} \frac{1}{2\sqrt{y}} + \frac{1}{\sqrt{2\pi}} e^{-\frac{(-\sqrt{y})^2}{2}} \frac{1}{2\sqrt{y}} = \frac{y^{-1/2} e^{-\frac{y}{2}}}{\sqrt{\pi}} \end{aligned}$$

Compare with the PDF of Gamma RV ($f(y) = \frac{y^{\alpha-1} e^{-\frac{y}{\beta}}}{2^{1/2} \beta^{\alpha} \Gamma(\alpha)}$) gives $\alpha = 1/2$, $\beta = 2$, $\Gamma(1/2) = \sqrt{\pi}$

Normal vs χ^2

Z standard normal RV $\implies Z^2$ is Gamma RV with $\alpha = \frac{1}{2}, \beta = 2$ ($=\chi^2$ RV)

Simulation of Random Variable on a computer

The only source of randomness on a computer is a (pseudo-) number generator which gives a uniform random variable U on $[0, 1]$. How do we generate other random variables?

The inverse CDF method

Suppose Y is a continuous random variable with CDF $F(y)$ and U is uniform on $[0, 1]$. Then we have

$$Y = F^{-1}(U)$$

Proof We check that $Y = F^{-1}(U)$ has the correct CDF, namely F itself.

$$\begin{aligned} P(Y \leq y) &= P(F^{-1}(U) \leq y) \\ &= P(U \leq F(y)) \quad (F \text{ invertible}) \\ &= F(y) \quad (\text{since } P(U \leq a) = a) \end{aligned}$$

Example An exponential random variable with parameter β has CDF $F(y) = 1 - e^{-y/\beta}$. The inverse function F^{-1} is

$$z = 1 - e^{-y/\beta} \Leftrightarrow e^{-y/\beta} = 1 - z \Leftrightarrow y = -\beta \ln(1 - z)$$

So the inverse function of $F(y) = 1 - e^{-y/\beta}$ is $F^{-1}(z) = -\beta \ln(1 - z)$ and so

$$Y = -\beta \ln(1 - U)$$

has an exponential distribution with parameter β .

Pseudocode to generate exponentials random variables

- Generate a random number U .
- Set $Y = -\beta \ln(1 - U)$

The method of transformation

This is **special case of the CDF method which avoids the computation of integral**

The transformation method

- The RV Y has pdf $f_Y(y)$.
- For the RV $Z = h(Y)$ we assume that h is an increasing (or decreasing function)

if $y_1 < y_2$ then $h(y_1) < h(y_2)$ (or $h(y_2) < h(y_1)$).

and so the inverse function $h^{-1}(z)$ exists.

- Then the RV Z has pdf

$$f_Z(z) = f_Y(h^{-1}(z)) \left| \frac{dh^{-1}(z)}{dz} \right|$$

Example for the transformation method

- Suppose Y is an exponential random variable, we want to find the distribution of $Z = \sqrt{Y}$.
- The transformation $h(y) = \sqrt{y}$ is invertible with inverse

$$h(y) = \sqrt{y} = z \iff y = z^2 = h^{-1}(z)$$

- The derivative is

$$\frac{d}{dz}h^{-1}(z) = \frac{d}{dz}z^2 = 2z$$

- The density of Z is

$$f_Z(z) = f_Y(h^{-1}(z)) \left| \frac{dh^{-1}(z)}{dz} \right| = z f_Y(z^2) = \frac{z}{\beta} e^{-z^2/\beta}$$

Z is called a Weibull random variable.

More examples: functions of 2 uniform random variables

X_1 and X_2 independent and uniform random variables on $[0, 1]$ so the joint PDF is

$$f(x_1, x_2) = 1 \quad \begin{array}{l} 0 \leq x_1 \leq 1 \\ 0 \leq x_2 \leq 1 \end{array}$$



Since the density is constant we can compute

$$P((X_1, X_2) \in A) = \text{area of } A \quad (\text{maybe avoid integrals})$$

Example 1: Find the PDF of $Y = X_1 + X_2$.

We have $0 \leq Y = X_1 + X_2 \leq 2$ and we need to compute the CDF

$F_Y(y) = P(Y \leq y)$ for $0 \leq y \leq 2$. There are 2 cases.




$$P(Y \leq y) = \frac{y^2}{2}$$


$$P(Y \leq y) = 1 - \frac{(2-y)^2}{2}$$

So we have

$$\text{PDF } f(y) = F'(y) = \begin{cases} y & \text{if } 0 \leq y \leq 1 \\ 2-y & \text{if } 1 \leq y \leq 2 \end{cases}$$



Example 2: Find the PDF of $Y = X_1 X_2$. We have $0 \leq Y \leq 1$ and we need to draw the region $\{x_2 \leq \frac{y}{x_1}\}$.



$$P(Y \leq y) = \text{area A} + \text{area B}$$

$$\text{area A} = y$$

$$\text{area B} = \int_y^1 \int_0^{y/x_1} 1 dx_2 dx_1$$


$$= \int_y^1 \frac{y}{x_1} dx_1$$

$$= y \ln(x_1) \Big|_{x_1=y}^{x_1=1}$$

$$= -y \ln(y)$$

$$\text{So } F(y) = y - y \ln(y)$$

$$\text{PDF } f(y) = F'(y) = -\ln(y) \quad 0 \leq y \leq 1$$



Multivariable transformations

- The RV $Y = (Y_1, Y_2)$ has joint pdf $f_Y(y_1, y_2)$.
- Consider the random variable $Z = (Z_1, Z_2)$ given by
$$\begin{aligned}Z_1 &= h_1(Y_1, Y_2) \\ Z_2 &= h_2(Y_1, Y_2)\end{aligned}$$
- Assume the transformation is invertible
$$\begin{aligned}Y_1 &= h_1^{-1}(Z_1, Z_2) \\ Y_2 &= h_2^{-1}(Z_1, Z_2)\end{aligned}$$

The multivariate transformation method

The RV Z has pdf

$$f_Z(z) = f_Y(h^{-1}(z))J(z)$$

where the **Jacobian** is given

$$J(z) = \left| \det \begin{pmatrix} \frac{\partial h_1^{-1}}{\partial z_1}(z_1, z_2) & \frac{\partial h_1^{-1}}{\partial z_2}(z_1, z_2) \\ \frac{\partial h_2^{-1}}{\partial z_1}(z_1, z_2) & \frac{\partial h_2^{-1}}{\partial z_2}(z_1, z_2) \end{pmatrix} \right|$$

Recall that $\det \begin{pmatrix} a & b \\ c & d \end{pmatrix} = ad - bc$

Polar coordinate and Box Muller

Polar coordinates:
$$\begin{aligned} r &= \sqrt{x^2 + y^2} \\ \theta &= \tan^{-1}(y/x) \end{aligned} \iff \begin{aligned} x &= r \cos(\theta) \\ y &= r \sin(\theta) \end{aligned}$$

The **Jacobian** is

$$J(r, \theta) = \left| \det \begin{pmatrix} \cos(\theta) & \sin(\theta) \\ -r \sin(\theta) & r \cos(\theta) \end{pmatrix} \right| = r(\cos^2(\theta) + \sin^2(\theta)) = r$$

Suppose (X, Y) are independent standard normal with joint pdf

$$f(x, y) = \frac{e^{-x^2/2}}{\sqrt{2\pi}} \frac{e^{-y^2/2}}{\sqrt{2\pi}}$$

then the joint density of (R, Θ) is

$$f_{R, \Theta}(r, \theta) = f_{X, Y}(r \cos(\theta), r \sin(\theta)) r = r e^{-r^2/2} \frac{1}{2\pi}$$

Θ and R are independent and Θ is uniform on $[0, 2\pi]$ and R has density $re^{-r^2/2}$ on $[0, \infty)$.

The CDF for R is $P(R \leq r) = F(r) = 1 - e^{-r^2}$.

The inverse function is

$$1 - e^{-r^2} = u \iff u = \sqrt{-\ln(1 - z)}$$

so by the inverse CDF methods we have

$$R = \sqrt{-\ln(1 - U)}$$

Box-Muller Algorithm:

To generate two independent standard normal X, Y use two random number and set

$$X = \sqrt{-\ln(1 - U_1)} \cos(2\pi U_2)$$

$$Y = \sqrt{-\ln(1 - U_1)} \sin(2\pi U_2)$$